

*bancrofti*

*Wuchereria*

F. O. RICHARDS JR\*, D. D. PAM†, A. KAL, G. Y. GERLONG, J. ONYEKA†, Y. SAMBO, J. DANBOYI§, B. IBRAHIM¶, A. TERRANELLA\*, D. KUMBAK†, A. DAKUL†, A. LENHART\*, L. RAKERS\*, J. UMARU‡, S. AMADIEGWU, P. C. WITHERS JR\*, H. MAFUYAI†, M. Y. JINADU\*\*, E. S. MIRI and A. EIGEGE

\*T C C, O C, 453 F P, A, GA 30307, U.S.A  
†U ♀, D, P.M.B. 2084, ♀ P S, N  
‡T C C /G 2000, 1♀ K S, O T W R R, P.O. B / 7772, ♀, P S, N  
§S M H, P.M.B. 032, L, N S, N  
¶S M H, N P I (NPI), P 1266 A  
B W, A II, P.M.B. 511, G, A, N  
\*\*F M H, L O, 9 F P II, F S, I - L, N  
R 1 S 2004, R 7 O 2004,  
A 11 O 2004

A prospective entomological survey was conducted in four sentinel villages in central Nigeria from 1999–2002, to assess the impact of annual, single-dose, mass drug administrations (MDA), with a combination of ivermectin and albendazole, on the transmission of



zones (unpubl. obs.). The principal local

could be sampled during each morning's





Maiganga, Lankan and Gbuwhen were 82%, 78% and 63%, respectively.

## DISCUSSION

The high prevalences of infection and infectivity recorded in the mosquitoes at the start of the present study indicate that 5–7 years of annual mass treatments with ivermectin alone failed to interrupt transmission of *W. bancrofti* in the study area, where *A. gambiae* s.l. appears to be the main vector. In Burkina Faso, Kyelem *et al.* (2003) similarly reported that 5 years of semi-annual mass treatments with ivermectin alone (targetted at onchocerciasis) reduced but did not interrupt the transmission of *W. bancrofti*.

During the present study, the addition of albendazole to the ivermectin-based MDA led, after three rounds of treatment, to significant reductions in the prevalences of mosquito infection in three of the four sentinel villages. Annual mass treatment with ivermectin–albendazole therefore appears superior to annual ivermectin monotherapy for lowering or stopping the transmission of *W. bancrofti*, at least in rural areas of sub-Saharan Africa. The present results appear to represent the first entomological confirmation of the importance of albendazole in MDA-based programmes of LF control (Addiss *et al.*, 1997, 2004; Ismail *et al.*, 1998; Ottesen *et al.*, 1999; Dunyo *et al.*, 2000), although the present study has some limitations that need to be considered. Firstly, as there were no human-landing catches, it was impossible to determine

highest interval treatment coverage (86%) of the four sentinel villages. There was, however, no evidence of the mosquito infections that were detected clustering by compounds in any of the four villages (data not shown), indicating that the present data were not influenced by a few infected villagers who repeatedly refused treatment.

As the prevalence of mosquito infection is gradually reduced by repetitive rounds of MDA, entomological-impact monitoring (at least that based on traditional dissection methods) will become ever more difficult,



## REFERENCES

- Addiss, D., Critchley, J., Ejere, H., Garner, P., Gelband, H. & Gamble, C. (2004). Albendazole for lymphatic filariasis (Cochrane Review). In *The Cochrane Library*; Issue 4, 2004. Chichester, U.K.: John Wiley & Sons.
- Addiss, D. G., Beach, M. J., Streit, T. G., Lutwick, S., LeConte, F. H., Lafontant, J. G., Hightower, A. W. & Lammie, P. J. (1997). Randomised placebo-controlled comparison of ivermectin and albendazole alone and in combination for *Wuchereria bancrofti* microfilaraemia in Haitian children. *Lancet*, **350**, 480-484.
- Awadzi, K., Edwards, G., Duke, B. O. L., Opoku, N. O., Attah, S. K., Addy, E. T., Ardrey, A. E. & Quartey, B. T. (2003). The co-administration of ivermectin and albendazole — safety, pharmacokinetics and efficacy against *Onchocerca volvulus*. *Acta Tropica*, **97**, 165-178.
- Bockarie, M. J., Tisch, D. J., Kastens, W., Alexander, N. D., Dimber, Z., Bockarie, F., Ibam, E., Alpers, M. P. & Kazura, J. W. (2002). Mass treatment to eliminate filariasis in Papua New Guinea. *New England Journal of Medicine*, **347**, 1841-1848.
- Brown, K. R., Ricci, F. M. & Ottesen, E. A. (2000). Ivermectin: effectiveness in lymphatic filariasis. *Parasitology*, **121**, 133-146.
- Burnham, G. (1998). Onchocerciasis. *Lancet*, **351**, 1341-1346.
- Dunyo, S. K., Nkrumah, F. K. & Simonsen, P. E. (2000). Single-dose treatment of *Wuchereria bancrofti* infections with ivermectin and albendazole alone or in combination: evaluation of the potential for control at 12 months after treatment. *Tropical Medicine and International Health*, **94**, 437-443.
- Eigege, A., Richards, F. O., Blaney, D. D., Miri, E. S., Gontor, I., Ogah, G., Umaru, J., Jinadu, M. Y., Mathai, W., Amadiogwu, S. & Hopkins, D. R. (2003). Rapid assessment for lymphatic filariasis in central Nigeria: a comparison of the immunochromatographic card test and hydrocele rates in an area of high endemicity. *Acta Tropica*, **68**, 643-646.

- treatment for lymphatic filariasis and onchocerciasis. *T P*, **19**, 516-522.
- Ottesen, E. A., Duke, B. O., Karam, M. & Behbehani, K. (1997). Strategies and tools for the control/elimination of lymphatic filariasis. *B W H O*, **75**, 491-503.
- Ottesen, E. A., Ismail, M. M. & Horton, J. (1999). The role of albendazole in programmes to eliminate lymphatic filariasis. *P T*, **15**, 382-386.
- Plaisier, A. P., Stolk, W. A., van Oortmarssen, G. J. & Habbema, J. D. (2000). Effectiveness of annual ivermectin treatment for *W* infection. *P T*, **16**, 298-302.
- Richards, F., Gonzales-Peralta, C., Jallah, E. & Miri, E. (1996). Community-based ivermectin distributors: onchocerciasis control at the village level in Plateau state, Nigeria. *A T*, **61**, 137-144.
- Richards, F., Hopkins, D. & Cupp, E. (2000). Programmatic goals and approaches to onchocerciasis. *L*, **355**, 1663-1664.
- Weil, G. J., Lammie, P. J. & Weiss, N. (1997). The ICT Filariasis test: a rapid-format antigen test for diagnosis of bancroftian filariasis. *P T*, **13**, 401-404.
- Williams, S. A., Laney, S. J., Bierwert, L. A., Saunders, L. J., Boakye, D. A., Fischer, P., Goodman, D., Helmy, H., Hoti, S. L., Vasuki, V., Lammie, P. J., Plichart, C., Ramzy, R. M. & Ottesen, E. A. (2002). Development and standardization of a rapid, PCR-based method for the detection of *W* in mosquitoes, for xenomonitoring the human prevalence of bancroftian filariasis. *A T M P*, **96** (Suppl. 2), S41-S46.
- World Health Organization (1995). *O C R WHO E C O C*. Technical Report Series No. 852. Geneva: WHO.
- World Health Organization (2002). *D P E L F R I C WHO/HQ, G*, 29 31 ¶ 2002. Document WHO/CDS/CPE/PVC/2002.3. Geneva: WHO.